

Influence of artificial and native extracellular matrices on adhesion and differentiation of mesenchymal stem cells and osteoblasts

[Dr. rer. nat. Ute Hempel](#) , Sarah Vogel, Carolin Preißler (MTA)

Artificial extracellular matrices (aECM) consisting of collagen and sulfated glycosaminoglycan derivatives (sGAG) promote the osteogenic lineage commitment of human mesenchymal stromal cells (hMSC). This effect was seen with sGAG in 2D (substrate coatings) and in 3D (modified fibrillary collagen networks, B10).

In addition to the pro-osteogenic effects, sGAG and aECM alter significantly the endogenous (native) ECM (nECM) of hMSC. sGAG derivatives interact with hMSC-derived matrix vesicles, they co-localize with numerous nECM proteins (e.g. fibronectin) and become incorporated in the complex nECM network. Thus, sGAG induce changes in the nECM composition, assembly, cross-linking, remodelling, and mineralisation (cooperation with Z4 (proteomics), A3 (SPR analysis), A7 (molecular modelling), and Viola Vogel's group, ETH Zurich FRET, fiber stretch assay).

For better understanding how EZM characteristics influence osteogenic differentiation we will investigate sGAG-induced alterations of nECM structure, composition and physico-chemical properties as well as the phosphorylation status of particular ECM proteins more detailed (cooperation with A3, A6, B10, Z4). Our working hypothesis is that the pro-osteogenic effect of sGAG derivatives and aECM is mainly caused by sGAG-induced altered nECM. Therefore, the so altered nECM itself should be used for the modification of biomaterial surfaces. Our studies showed further an intracellular enrichment of sGAG and suggested that the endocytotic receptor LRP-1/CD91 (low density lipoprotein receptor-related protein-1) could be involved. Pull-down experiments and SPR analyses (cooperation with Z4 and A3) are planned to identify intracellular target molecules.

According to the strategy of the consortium we investigate whether sGAG and ECM reveal beneficial effects for hMSC from different sources and patients with compromised bone healing (cooperation with B5 and the Savkovic group).





Figure 1. Collagen | F-Actin Morphology of hMSC on (left) and in (right) a 3D collagen network, 24 h after plating

Project-related Publications

1. Rother S, Galiazzo V, Kilian D, Fiebig K, Becher J, Möller S, [Hempel U](#), [Schnabelrauch M](#), Waltenberger J, [Scharnweber D](#)

, [Hintze V](#)

. Hyaluronan/collagen hydrogels with sulfated hyaluronan for improved repair of vascularized tissue tune the binding of proteins and promote endothelial cell growth.

Macromolecular Bioscience.

2017;

accepted

2. Köhler L, Samsonov S, Rother S, Vogel S; Köhling S, Möller S, [Schnabelrauch M](#), [Rademann J](#)

, [Hempel U](#)

, [Pisabarro MT](#)

, [Scharnweber D](#)

, [Hintze V](#)

. Sulfated hyaluronan derivatives modulate TGF- β 1:receptor complex formation: Possible consequences for TGF- β 1 signaling.

[Scientific Reports.](#)

[2017;7:1210.](#)

3. Vogel S, Arnoldini S, Möller S, Schnabelrauch M, [Hempel U](#). Sulfated hyaluronan alters fibron-ectin matrix assembly and promotes osteogenic differentiation of human bone marrow stromal cells. [Sci Rep.](#)

[2016;6:36418.](#)

4. Schmidt J, Kliemt S, Preissler C, Möller S, von Bergen M, [Hempel U*](#), Kalkhof S*. Osteoblast-released matrix vesicles - Regulation of activity and composition by sulfated and non-sulfated glycosaminoglycans.

[Mol Cell Proteomics.](#)

[2016;15:558-572.](#)

(*geteilte Seniorautorschaft)

5. [Hempel U](#), Müller K, Preissler C, Noack C, Boxberger S, Dieter P, Bornhäuser M, Wobus M. Human bone marrow stromal cells: A reliable, challenging tool for in vitro osteogenesis and bone engineering approaches. [Stem Cells Int .](#)

[2016;7842191](#)

6. Lutter AH, [Hempel U](#), Anderer U, Dieter P. Biphasic influence of PGE2 on the resorption activity of osteoclast-like cells derived from human peripheral blood monocytes and mouse RAW264.7 cells. [PLEFA . 2016;111:1-7 .](#)

7. Rother S, Samsonov S, [Hempel U](#), Vogel S, Möller S, Blaszkiewicz J, Köhling S, Schnabelrauch M, Rademann J, Pisabarro MT, Hintze V, Scharnweber, D. Sulfated Hyaluronan Alters the Interaction Profile of TIMP-3 with the Endocytic Receptor LRP-1

Clusters II and IV and Increases the Extracellular TIMP-3 Level of Human Bone Marrow Stromal Cells.

[Biomacromolecules](#) .

[2016;17:3252-61.](#)

8. Noack C, [Hempel U](#), Preissler C, Dieter P. Prostaglandin E2 impairs osteogenic and facilitates adipogenic differentiation of human mesenchymal stromal cells.

[Prostaglandins, Leukotrienes & Essential Fatty Acids](#)

[2015;94:91-98.](#)

9. Elschner C, Noack C, Preißler C, Krause A, Scheler U, [Hempel U](#). In vitro response of human mesenchymal stem cells to titanium coated PEEK films and their suitability for magnetic resonance imaging.

[Journal of Materials Science & Technology](#)

[2015;5:427-436](#)

10. Miron A, Rother S, Huebner L, [Hempel U](#), Käßler I, Moeller S, Schnabelrauch M, Scharnweber D, Hintze V. Sulfated hyaluronan influences the formation of artificial extracellular matrices and the adhesion of osteogenic cells.

[Macromol Biosci](#)

[. 2014 Dec;14\(12\):1783-94. doi: 10.1002/mabi.201400292. Epub 2014 Sep 12](#)

11. Kalkhof S, Förster Y, Schmidt J, Schulz M, Baumann S, Weißflog A, Gao W, [Hempel U](#), Eckelt U, Rammelt S, Von Bergen M. Proteomics and metabolomics for in situ monitoring of wound healing.

[Biomed Res Int](#)

[. 2014;2014:934848.](#)

12. [Hempel U](#), Preissler C, Vogel S, Möller S, Hintze V, Becher J, Schnabelrauch M, Rauner M, Hofbauer LC, Dieter P. Artificial extracellular matrices with oversulfated glycosaminoglycan derivatives promote the differentiation of osteoblast-precursor cells and premature osteoblasts.

[J Biomed Res Int](#) [2014;938368](#) .

13. [Hempel U](#), Matthäus C, Preissler C, Möller S, Hintze V, Dieter P. Artificial matrices with high-sulfated glycosaminoglycans and collagen are anti-inflammatory and pro-osteogenic for human mesenchymal stromal cells.

[J Cell Biochem](#) [2014,115:1561-1571](#) .

14. Mai M, [Hempel U](#), Hacker M, Dieter P. Effects of HyStem-HP™ hydrogel elasticity on osteogenic differentiation of human mesenchymal stromal cells.

[Cell Mol Bioneng](#)

[2014;7:155-164.](#)

15. Förster Y, Gao W, Demmrich A, [Hempel U](#), Hofbauer LC, Rammelt S. Monitoring the first stages of bone healing with microdialysis.

[Acta Orthopaedica](#)

[2013;84\(1\):76-81](#)

16. Kliemt S, Lange C, Otto W, Hintze V, Möller S, von Bergen M, [Hempel U](#), Kalkhof S. Sulfated hyaluronan containing collagen matrices enhance cell-matrix-interaction, endocytosis, and osteogenic differentiation of human mesenchymal stromal cells.

[J Proteome Res](#)

[. 2013;12:378-89](#)

17. Büttner M, Möller S, Keller M, Huster D, Schiller J, Schnabelrauch M, Dieter P, [Hempel](#)

U

Over-sulfated chondroitin sulfate derivatives induce osteogenic differentiation of human mesenchymal stromal cells independent of bone morphogenetic protein-2 and transforming growth factor- β 1 signalling.

[J Cell Physiology](#)
[. 2013;228:330-40](#)

18. Hempel U, Möller S, Noack C, Hintze V, Scharnweber S, Schnabelrauch M, Dieter P. Sulfated hyaluronan/collagen I-matrices enhance osteogenic differentiation of human mesenchymal stromal cells in vitro even in the absence of dexamethasone.

[Acta Biomater](#)
[. 2012;8:4064-72](#)

19. Hempel U, Hintze V, Möller S, Schnabelrauch M, Scharnweber D, Dieter P. Artificial extracellular matrices composed of collagen I and sulfated hyaluronan with adsorbed TGF- β 1 promote collagen synthesis of human mesenchymal stromal cells. [Acta](#)

[Biomater](#)
[. 2012;8:659-6](#)

6.

20. Salbach J, Rachner T, Rauner M, Hempel U, Anderegg U, Franz S, Simon JC, Hofbauer LC. Regenerative potential of glycosaminoglycans for skin and bone.

[J Mol Med](#)
[. 2012;90:620-35](#)

21. Hempel U, Hintze V, Möller S, Schnabelrauch M, Scharnweber D, Dieter P. Artificial extracellular matrices composed of collagen I and sulfated hyaluronan with adsorbed transforming growth factor β 1 promote collagen synthesis of human mesenchymal stromal cells. [Acta Biomater](#) . 2012;8:659-66.

22. Hamann C, Götsch C, Mettelsiefen J, Henkenjohann V, Rauner M, Hempel U, Bernhardt R, Fratzl-Zelman N, Roschger P, Rammelt S, Günther KP, Hofbauer LC. Delayed bone regeneration and low bone mass in a rat model of insulin-resistance type 2 diabetes mellitus is due to impaired osteoblast function.

[Am J Physiol Endocrinol Metab.](#)
[2011;301:e1220-8](#)

23. Goettsch C, Rauner M, Pacyna M, Hempel U, Bornstein SR, Hofbauer LC. mi-R-125b regulates calcification of vascular smooth muscle cells.

[Am J Pathol](#)
[. 2011;179:1594-600](#)

.Goettsch C, Rauner M, Hamann C, Sinnigen K,
Hempel U

, Bornstein SR, Hofbauer LC. Nuclear factor of activated T cells mediates oxidised LDL-induced calcification of vascular smooth muscle cells.

[Diabetologia](#)
[. 2011;54:2690-701](#)

24. Kunze R, Hempel U, Dieter P. Differential Effect of Chondroitin-4-sulfate on the immediate and delayed Prostaglandin E(2) Release from Osteoblasts.

[Prostaglandins other Lipid Mediat](#)

[. 2010;epub](#)

25. Lutter AH, Hempel U, Wolf-Brandstetter C, Garbe A, Goettsch C, Hofbauer LC, Jessberger R, Dieter P. A novel resorption assay for osteoclast functionality based on an osteoblast-derived native extracellular matrix.

[J Cell Biochem](#)

[. 2010;109:1025-32](#)

26. Kunze R, Rösler M, Möller S, Schnabelrauch M, Riemer T, Hempel U, Dieter P. Sulfated hyaluronan derivatives reduce the proliferation rate of primary rat calvarial osteoblasts.

[Glycoconj](#)

[. 2010;27:151-8](#)

27. Helas S, Goettsch C, Schoppet M, Zeitz U, Hempel U, Morawietz H, Kostenuik PJ, Erben RG, Hofbauer LC. Inhibition of Receptor Activator of NF-kappa B Ligand by Denosumab Attenuates Vascular Calcium Deposition in Mice.

[Am J Pathol](#)

[. 2009;175:2249.](#)

28. Scheffler A-H, Hempel U, Dieter P. Verfahren und Mittel zum Nachweis der Aktivität von Osteoklasten. (Deutsche Patentanmeldung Nr. 10 2009 013 957.5-41)